

***Trirhithrum culcasiae* White & Copeland**

Trirhithrum culcasiae White & Copeland, 2003: 92.

Wing length=2.8-4.0 mm; Aculeus length=0.68 mm.

Male

Head: Arista long plumose. Two pairs frontal setae. Face black except for pale lateral and dorsal margins.

Thorax: Postpronotal lobe with a dark central mark. Scutum with thin microtrichose covering. Scutellum disk largely dark and marked with a narrow wavy yellow transverse line. Anepisternum largely dark; dorsal quarter pale; one seta.

Wing: Pattern distinct. Subbasal and discal crossbands indistinctly separated by numerous hyaline flecks; cell c extensively hyaline; discal crossband beyond pterostigma, and R-M crossvein within discal crossband. Subapical crossband joined to discal crossband; base deep, partly in cell dm. Posterior apical crossband reduced to a short spur. Anal lobe coloured but with a hyaline indentation (ending before vein A₁+Cu₂). No bulla.

Legs: Femora dark.

Abdomen: With reticulate grey/silvery microtrichose pattern.

Female

Terminalia: Aculeus fairly short and pointed (appears asymmetric under a coverslip; dorsal view apparently similar to *T. leonense*; spermatheca with a bulbous apex, apparently similar to *T. occipitale*. (description after White et al., 2003)