

***Trirhithrum fraternum* Munro**

[*Ceratitis*] *Trirhithrum fraternum* Munro, 1934: 482

Wing length=3.7-4.2 mm; Aculeus length=0.94 mm.

Male

Head: Arista plumose. Two pairs frontal setae. Face white.

Thorax: Postpronotal lobe entirely dark or sometimes with an indistinct pale margin. Scutum without silvery-white microtrichose areas. Scutellum disk dark; margin with baso-lateral pale spots; spots adjacent to bases of apical setae. Anepisternum entirely dark; one seta. Anatergite without a bright silvery spot.

Wing: Pattern distinct. Subbasal and discal crossbands fused posterior to Rs and cell c extensively hyaline; cell bc with dark area not extended into basal half of cell. Discal crossband distally aligned with a point within pterostigma and R-M crossvein within discal crossband. Subapical crossband not joined to discal crossband (rarely almost joined). Posterior apical crossband reduced to a short spur. Anal lobe coloured but with a hyaline indentation (ending before vein  $A_1+Cu_2$ ). No bulla.

Legs: Femora dark.

Abdomen: With distinct grey microtrichose stripes.

Female

Terminalia: Aculeus fairly short and pointed (appears asymmetric when viewed under a coverslip; dorsal view apparently similar to *T. leonense*); spermatheca curved and bulbous (similar to *T. occipitale*). (description after White et al., 2003)