

***Trirhithrum quadrimaculatum* White**

Trirhithrum quadrimaculatum White, 2003: 115.

Wing length=3.9-4.5 mm; Aculeus length=1.08 mm.

Male

Head: Arista long pubescent to plumose. Two pairs frontal setae. Face entirely white.

Thorax: Postpronotal lobe pale either along lateral margin, or around entire margin. Scutum without silvery-white microtrichose areas. Scutellum disk dark; margin with baso-lateral pale spots; spots adjacent to bases of apical setae. Anepisternum largely dark; dorsal edge narrowly pale; one seta. Anatergite without a bright silvery spot. Katatergite with a small pale spot.

Wing: Pattern distinct. Subbasal and discal crossbands fused posteriorly and cell c extensively hyaline; cell bc with dark area not extended into basal half of cell. Discal crossband distally aligned with a point within pterostigma and R-M crossvein within discal crossband. Subapical crossband joined to discal crossband; base narrow, largely or entirely confined to cell r_{4+5} . Posterior apical crossband present to beyond vein M but not reaching wing margin. Anal lobe coloured but with a hyaline indentation (ending before vein A_1+Cu_2). No bulla. Legs: Femora dark.

Abdomen: With distinct grey microtrichose spots which tend to coalesce into stripes.

Female

Terminalia: Aculeus short, stout and apically pointed (does not appear asymmetric under a coverslip indicating that it is dorso-ventrally flattened); spermatheca apically bulbous.

(description after White et al., 2003)