

***Trirhithrum stecki* White**

Trirhithrum stecki White, 2003: 119.

Wing length=4.6-7.4 mm; Aculeus length=2.02 mm.

Male

Head: Arista long pubescent to plumose. Two pairs frontal setae. Face pale.

Thorax: Postpronotal lobe entirely dark. Scutum without silvery-white microtrichose areas. Scutellum disc white; margin dark between setae. Anepisternum with a diagonal white stripe covering posterior half; one seta. Anatergite without a bright silvery spot.

Wing: Pattern distinct. Subbasal and discal crossbands fused posteriorly and cell c extensively hyaline; discal crossband distally aligned with a point within pterostigma and R-M crossvein aligned to edge of discal crossband. Subapical crossband joined to discal crossband; base deep, partly in cell dm. Posterior apical crossband represented by both a spur off the costal band and by a separate (isolated) diagonal line which crosses vein M and more or less reaches the wing margin. Anal lobe largely dark, at most tending to hyaline along wing margin. No bulla.

Legs: Femora dark.

Abdomen: With grey/silvery microtrichose bands (not very distinct).

Female

Terminalia: Aculeus fairly long and pointed; spermatheca large, curved and slightly bulbous at apex (similar to *T. validum*).

(description after White et al., 2003)