

Dacus (Dacus) chrysomphalus (Bezzi)

Tridacus chrysomphalus Bezzi, 1924b: 453

Wing length, 7.3-7.7 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present (lateral fairly short, barely extending beyond anterior supra-alar seta). Scutellum without any dark patterning (except for basal dark margin, which is deep). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein R_{4+5} before wing apex; apically expanded into a spot which reaches vein M and starts before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu). Cell bc hyaline; cell c slightly coloured (almost as deep as costal band). Without any crossbanding.

Legs. Femora bicoloured (pale basally and red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)