

***Bactrocera (Bactrocera) dorsalis* (Hendel)**

*Dacus dorsalis* Hendel, 1912: 18

Wing length, 5.42-6.92 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum red-brown to black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vitta present; medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present; prescutellar acrostichal seta present; basal scutellar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not or only slightly extending below vein  $R_{2+3}$  before wing apex; not expanded into an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale coloured. Abdomen. Predominantly fulvous; shape and patterning, see image. Tergites II-V separate. Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae. Female. Aculeus pointed; no torsion; length, 1.3-1.8mm.

(Description after White, 2006)