

Bactrocera (Daculus) biguttula (Bezzi)

Chaetodacus biguttulus Bezzi, 1922: 294

Wing length, 4.2-6.2 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red brown, with a black stripe; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vittae present (lateral fairly short, barely extending beyond anterior supra-alar seta); medial postsutural vitta absent. Scutellum with a broad red-brown stripe from base to apex (lateral margin yellow). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent; prescutellar acrostichal seta present; basal scutellar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with microtrichia confined to anteriorly half. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; barely expanded at apex. Anal streak present but narrow. Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale; rarely with mid- and hindfemora bicoloured (pale basally, red-brown apically). Abdomen. Red-brown, patterned fuscous; shape and patterning, see image. Tergites II-V separate.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion; length, 1.04mm.

(Description after White, 2006)