

***Bactrocera (Daculus) lucida* (Munro)**

*Dacus (Afrodacus) lucidus* Munro, 1939a: 26

Wing length, 5.1-5.7 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 1, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus sometimes yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum concolorous with scutum. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; usually extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent; prescutellar acrostichal seta present; basal scutellar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent (reduced to a small apical spot). Anal streak present but narrow. Cells bc and c hyaline. Crossbanding; isolated crossband on R-M.

Legs. Femora pale coloured.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites II-V separate.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)