

***Bactrocera (Daculus) menanus* (Munro)**

*Afrodacus menanus* Munro, 1984: 24

Wing length, 5.3 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, with a small red (HT B. *menanus*) to black spot (HT B. *andriae*) on mesal side of antennal furrow. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vitta present; medial postsutural vitta absent. Scutellum with a narrow apical red-brown marking. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined (or almost so) to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent; prescutellar acrostichal seta present; basal scutellar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent; apical spot very large, contacting or crossing vein M and starting before end of vein R<sub>2+3</sub>. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basally and red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites II-V separate.

Male. Tergite III with pecten, dense microtrichia adjacent end A<sub>1</sub>+Cu<sub>2</sub> [hindtibia broken]. Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)