

***Bactrocera (Daculus) montyanus* (Munro)**

Mauritidacus montyanus Munro, 1984: 25

Wing length, 4.2-5.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vitta present; medial postsutural vitta absent. Scutellum with an apical black marking, or broad dark stripe from base to apex. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present or absent; prescutellar acrostichal seta present; basal scutellar seta present or absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent; apical spot very large, contacting or crossing vein M and starting before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale, sometimes with a dark prespaical marking, or bicoloured (pale basally, red-brown to fuscous apically).

Abdomen. Red-brown, patterned fuscous (including tergite III); shape and patterning, see image (CD-C). Tergites II-V separate.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)