

***Dacus (Lophodacus) chamun (Munro)***

*Nebrodacus chamun* Munro, 1984: 141

Wing length, 3.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 1-2, orbital setae 0.

Thorax. Scutum predominantly fulvous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; not extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with microtrichia confined to anteriorly half. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$ . Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal section with slightly thickened setae.

Female. Unknown.

(Description after White, 2006)