

***Dacus (Psilodacus) amberiens* (Munro)**

*Andriadacus amberiens* Munro, 1984: 70

Wing length, 5.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein  $R_{4+5}$  before wing apex; apically expanded into a spot which reaches vein M and starts before end of vein  $R_{2+3}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)