

***Dacus (Leptoxyda) apectus* White, 2006**

*Dacus (Leptoxyda) apectus* White, 2006: 98.

Wing length, 4.3-4.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, deep black band across lower two-thirds (no separate antennal furrow spot). Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured (largely yellow); notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent.

Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; narrowly extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; expanded slightly across apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Forefemur pale with a preapical dark marking; mid- and hindfemora bicoloured (midfemur pale in basal third, hindfemur pale basal two-thirds, red-brown to fuscous apically).

Abdomen. Red-brown, patterned black, including medial stripe on tergite IV; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)