

***Dacus (Psilodacus) herensis* (Munro)**

*Andriadacus herensis* Munro, 1984: 70

Wing length, 5.4 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline.

Without any crossbanding.

Legs. Femora bicoloured (pale basally and red-brown apically).

Abdomen. Predominantly red-brown [badly stained]; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae. [Female unknown.]

(Description after White, 2006)