

***Dacus (Psilodacus) hapalus* (Munro)**

Pycnodacus hapalus Munro, 1984: 133

Wing length, 5.2 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area dark. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus concolorous with scutum; notopleural xanthine apparently absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M before middle of cell dm (RMP=0.96). Costal band complete; shallow, not extending below vein R₂₊₃ before wing apex (except basally filling cell r₁); apically expanded into a spot which reaches about mid-depth of cell r₄₊₅. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding; crossband on R-M.

Legs. Femora pale coloured.

Abdomen. Predominantly black; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)