

***Dacus (Psilodacus) iaspideus* Munro**

Dacus (Psilodacus) iaspideus Munro, 1948b: 30

Wing length, 4.7-5.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area sometimes dark. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus sometimes with a small indistinct yellow spot; notopleural xanthine isolated from notopleural callus (or apparently absent); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; expanded into a small spot at apex. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)