

***Dacus (Psilodacus) inflatus* Munro**

*Dacus (Psilodacus) inflatus* Munro, 1939b: 3

Wing length, 4.6-5.3 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area sometimes dark. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus sometimes yellow; notopleural xanthine isolated from notopleural callus (or apparently absent); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present or absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band reduced to a very narrow band, often absent or barely discernible, and an apical spot. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding, when present, confined to a small isolated mark at upper end of R-M.

Legs. Femora pale coloured.

Abdomen. Predominantly black; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus apparently pointed and with preapical "shoulder" (not examined and no drawing available; described by Munro 1984: 133); no torsion.

(Description after White, 2006)