

***Dacus (Psilodacus) ruslan* (Hering)**

Psilodacus ruslan Hering, 1941b: 9

Wing length, 6.0-6.1 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area sometimes dark. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus concolorous with scutum; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R₄₊₅ before wing apex; expanded diagonally across apex to vein M. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding; crossband on R-M.

Legs. Femora bicoloured (forefemur indistinctly) (pale basally, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image. Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)