

***Dacus (Dacus) adenae* (Hering)**

Metidacus adenae Hering, 1940: 23

Wing length, 6.2-6.7 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum red-brown with dark stripes; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vittae present (lateral fairly short, barely extending beyond anterior supra-alar seta). Scutellum concolorous with scutum. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a separate xanthine on anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein R_{4+5} before wing apex; barely expanded at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)