

***Dacus (Dacus) bakingiliensis* Hancock**

Dacus (Dacus) bakingiliensis Hancock, 1985b: 303

Wing length, 4.9-6 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is deep). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly deep, extending below vein R_{2+3} or to vein R_{4+5} , before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding, when present, as a crossband along R-M.

Legs. Fore- and midfemora red-brown to fuscous, tending to bicoloured (pale basally, red-brown to fuscous apically); hindfemur red-brown (pale basally, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)