

***Dacus (Dacus) clinophlebs* Hendel**

Dacus clinophlebs Hendel, 1928: 344

Wing length, 5.2-5.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot or face almost entirely dark. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present (lateral fairly short, barely extending beyond anterior supra-alar seta). Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein R_{4+5} before wing apex; apically expanded into a spot which reaches below vein M; basally starting before R-M and extending in cell dm. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Fore- and midfemora bicoloured (pale basally, red-brown apically); hindfemur pale.

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)