

***Dacus (Dacus) collarti* Munro**

*Dacus (Dacus) collarti* Munro, 1938c: 163

Wing length, 5.3-5.7 mm.

*Head.* Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot (but very reduced in HT of *D. verecundus*). Frons, frontal setae 2, orbital setae 0.

*Thorax.* Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine largely confined to katatergite. *Thoracic setae.* Anterior notopleural seta present; anterior supra-alar seta present.

*Wing.* Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly deep, extending below vein  $R_{2+3}$  or to vein  $R_{4+5}$ , before wing apex; apically expanded into a spot which reaches nearly whole depth of cell  $r_{4+5}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

*Legs.* Femora pale coloured.

*Abdomen.* Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

*Male.* Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

*Female.* Unknown.

(Description after White, 2006)