

***Dacus (Dacus) croceus* Munro**

Dacus croceus Munro, 1957: 863

Wing length, 5.2-5.7 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with at most a very small dark spot. Frons, frontal setae 2, orbital setae 0-1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus with a small yellow spot; notopleural xanthine isolated from notopleural callus; lateral postsutural vitta absent; medial present. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an extensive covering of microtrichia (varies above and below 90%); cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R₄₊₅ before wing apex; apically expanded into a spot which extends almost to, or below, vein M, and starts before end of vein R₂₊₃. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (sometimes as deeply as costal band, especially cell c). Crossbanding; crossband on R-M.

Legs. Femora pale; forefemur sometimes with a preapical dark marking; midfemur sometimes bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A₁+Cu₂, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)