

***Dacus (Dacus) deltatus* White, 2006**

Dacus (Dacus) deltatus White, 2006: 66

Wing length, 4.6-4.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; entirely yellow. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured, or concolorous with scutum; notopleural callus concolorous with scutum, or at most yellow posteriorly; notopleural xanthine apparently absent (some suggestion of isolated xanthine in holotype); lateral postsutural vitta absent; medial variable (narrow trace in holotype; absent in MRAC paratype; well developed in TAU paratype). Scutellum with a basal black triangle or broad black stripe from base to apex. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; deep, extending to vein R_{4+5} before wing apex; apically expanded into a spot which reaches nearly whole depth of cell r_{4+5} and starts before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu).

Cells bc and c hyaline. Crossbanding; crossband on R-M triangular.

Legs. Femora pale; midfemur sometimes with a preapical dark marking.

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)