

Dacus (Dacus) disjunctus (Bezzi)

[*Dacus*] *Tridacus disjunctus* Bezzi, 1915: 96

Size. Medium, wing length, 6.9-7.3 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a separate xanthine on anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly shallow, slightly extending below vein R₂₊₃ before wing apex; not expanded into an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora bicoloured (pale basally, red-brown to fuscous apically); hindfemur pale.

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A₁+Cu₂, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)