

***Dacus (Dacus) famona* Hancock**

*Dacus (Ectopodacus) famona* Hancock, 1985b: 311

Wing length, 6.1-6.7 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is deep); sometimes concolorous with scutum. Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R<sub>4+5</sub> before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basally and red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A<sub>1</sub>+Cu<sub>2</sub>, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)