

***Dacus (Dacus) fumosus* Collart**

*Dacus (Dacus) fumosus* Collart, 1935a: 17

Wing length, 7.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a separate xanthine on anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M at or before middle of cell dm (RMP=0.92-1.0). Costal band complete; deep, extending to (or below) vein  $R_{4+5}$  before wing apex; apically expanded into a spot which reaches almost to vein M and starts before crossvein R-M. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Fore- and midfemora bicoloured (pale basally, red-brown apically); hindfemur pale.

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)