

***Dacus (Dacus) fuscinervis* Malloch**

Leptoxyda fuscinervis Malloch, 1932a: 301

Wing length, 6.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 0, orbital setae 0.

Thorax. Scutum predominantly red-brown; postpronotal lobe bicoloured; notopleural callus concolorous with scutum; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; not extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not or only slightly extending below vein R_{2+3} before wing apex (paler below vein R_{2+3} than above); barely expanded at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding; isolated crossbands on R-M and Dm-Cu.

Legs. Femora pale coloured. Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)