

***Dacus (Psilodacus) gabonensis* White, 2006**

*Dacus (psilodacus) gabonensis* White, 2006: 133

Wing length, 5.6 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper half of carina dark. Frons, frontal setae 1, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus with a small indistinct yellow spot; notopleural xanthine apparently absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Femora bicoloured; fore- and hindfemur pale basal three-quarters; midfemur pale basal two-thirds; red-brown apically.

Abdomen. Predominantly black; shape and patterning, see image; tergite IV with a medial stripe. Tergites I-V all fused.

Male. Unknown.

Female. No available data [tip broken].

(Description after White, 2006)