

***Dacus (Dacus) ghesquierei* Collart**

*Dacus (Dacus) ghesquierei* Collart, 1935a: 31

Wing length, 5.1 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot on a dark face. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly fuscous; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral postsutural vitta absent; medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent. Anal streak present (colour extending beyond cell bcu).

Cells bc and c hyaline. Crossbanding; isolated crossband on R-M.

Legs. Femora bicoloured (pale basally and red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)