

***Dacus (Dacus) hargreavesi* Munro**

Dacus (Dacus) hargreavesi Munro, 1939c: 139

Wing length, 5-5.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot or face almost entirely dark. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus (or apparently absent); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent (at most vein pale beyond cell sc). Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding; crossband on R-M.

Legs. Femora fuscous to black.

Abdomen. Predominantly black; tergite II largely black, or with an indistinct pale (sometimes divided) band, particularly in females; abdomen shape and other patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed, with 2 preapical "shoulders"; no torsion.

(Description after White, 2006)