

***Dacus (Dacus) ikelenge* Hancock**

*Dacus (Dacus) ikelenge* Hancock, 1985b: 300

Wing length, 7.5 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; not extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein  $R_{4+5}$  before wing apex; barely expanded at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and hindfemora bicoloured (pale basally, red-brown apically); midfemur red-brown.

Abdomen. Predominantly red-brown; shape and pattern, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ .

Female. Unknown.

(Description after White, 2006)