

***Dacus (Ambitidacus) katonae* Bezzi**

Dacus katonae Bezzi, 1924c: 90 [named but not fully described]

Wing length, 5.6-6.2 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; not extended onto katepisternum; anteriorly almost reaching postpronotal lobe. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; barely expanded at apex. Anal streak short but often reduced or absent. Cells bc and c hyaline. Crossbanding; crossband on R-M; often with a narrow crossband on DM-Cu.

Legs. Forefemur pale with a preapical dark marking; midfemur bicoloured (pale basally, fuscous apically); hindfemur pale.

Abdomen. Predominantly black; shape and patterning, see image. Tergites I-V all fused.

Male. Pecten absent [not verified from holotype which has a badly damaged abdomen]; no dense microtrichia area adjacent end A_1+Cu_2 [apparently absent in holotype]; no hindtibia preapical "pad".

Female. Aculeus pointed; no torsion.

(Description after White, 2006)