

***Dacus (Dacus) notalaxus* Munro**

*Dacus notalaxus* Munro, 1984: 39

Wing length, 6.2-6.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; carina black. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral postsutural vitta present; medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; very deep, extending below vein  $R_{4+5}$  before wing apex; not expanded into an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c slightly coloured (as deeply as costal band). Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten; lacking dense area of microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data [tip broken].

(Description after White, 2006)