

***Dacus (Dacus) pecropsis* Munro**

Dacus pecropsis Munro, 1984: 44

Wing length, 7.3-7.7 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a long dark stripe. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus sometimes with a small yellow spot; notopleural xanthine apparently absent (some suggestion of isolated xanthine in one specimen); lateral postsutural vitta sometimes present; medial absent. Scutellum concolorous with scutum. Anepisternum without a stripe; at most with an indistinct yellow spot. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present or absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete, pale between sc and darker apical spot; deep, extending to vein R₄₊₅ before wing apex; expanded into a small spot at apex; remainder of wing sometimes fumose. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A₁+Cu₂, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)