

***Dacus (Dacus) phimis* (Munro)**

Anomoiodacus phimis Munro, 1984: 59

Wing length, 4.9 mm.

Head. Pedicel+1st flagellomere length uncertain (head missing). Face, antennal furrow with a dark spot or face almost entirely dark. Frons, frontal setae 2, orbital setae 0-1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent (at most vein pale beyond cell sc). Anal streak present (colour extending beyond cell bcu). Cell bc hyaline; cell c slightly coloured (not as deep as costal band). Crossbanding; crossband on R-M.

Legs. Fore- and midfemora red-brown to black, tending to bicoloured (pale basally, red-brown to black apically); hindfemur bicoloured (pale basally, red-brown to black apically).

Abdomen. Predominantly black; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with some very fine hairs (presumed vestigial pecten); with barely differentiated area of microtrichia adjacent end A_1+Cu_2 , and slight hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)