

***Dacus (Ambitidacus) pulchralis* White, 2006**

Dacus (Ambitidacus) pulchralis White, 2006: 80

Wing length, 5.2-5.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; carina yellow; furrow orange-yellow. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band reduced to a very narrow band, often absent or barely discernable, and an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding; isolated crossbands on R-M and DM-Cu.

Legs. Femora pale coloured (almost white).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)