

***Dacus (Psilodacus) rutilus* Munro**

*Dacus (Dacus) rutilus* Munro, 1948b: 13

Wing length, 5.9 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus concolorous with scutum; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum concolorous with scutum. Anepisternum without a stripe. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band reduced to a very narrow band, often absent or barely discernable, and a small apical spot. Anal streak variable (sometimes confined to within bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora red-brown.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)