

***Dacus (Dacus) sakeji* Hancock**

*Dacus (Dacus) sakeji* Hancock, 1985b: 301

Wing length, 6.4 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum red-brown to black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is deep). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present or absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly deep, extending below vein  $R_{2+3}$  or to vein  $R_{4+5}$ , before wing apex; not expanded into an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding, when present, as a pale crossband along R-M.

Legs. Femora bicoloured (pale basally red-brown apically); midfemur tending to be almost entirely red-brown.

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)