

***Dacus (Psilodacus) schoutedeni* Collart**

*Dacus (Dacus) schoutedeni* Collart, 1935a: 11

Wing length, 5-5.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area dark especially in male. Frons, frontal setae 1, orbital setae 0-1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus with a small yellow spot; notopleural xanthine isolated from notopleural callus; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Forefemur pale; mid- and hindfemora bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, and hindtibia preapical "pad"; lacking dense area of microtrichia adjacent end  $A_1+Cu_2$ . Basal costal sections without specialised setae.

Female. Aculeus apex pointed (dorsal view); oblique truncate when viewed laterally; no torsion.

(Description after White, 2006)