

***Dacus (Didacus) bistrigulatus* Bezzi**

Dacus bistrigulatus Bezzi, 1908b: 193

Wing length, 4.6-5.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a long dark stripe. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. *Thoracic setae.* Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemora red-brown to bicoloured (pale basally, red-brown apically); all femora sometimes pale [probably only teneral specimens].

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus blunt, somewhat incised, and with preapical "shoulder"; complete torsion.

(Description after White, 2006)