

***Dacus (Didacus) briani* White, 2006**

Dacus (Didacus) briani White, 2006: 92

Wing length, 5.25 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous (dark red-brown with fuscous-black presutural and postsutural marks); postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta absent (a short and very thin white seta in/near the position on both sides of holotype); anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; barely expanded at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basal two-thirds to three-quarters, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)