

***Dacus (Didacus) congoensis* White, 2006**

Dacus (Didacus) congoensis White, 2006: 87

Wing length, 4.3-5.1 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot (paratype female) or face almost entirely dark (holotype male). Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band narrow, not extending below vein R_{2+3} before wing apex; at least one-third depth of cell r_{2+3} between end of vein R_{2+3} and apical spot; base of cell r_{2+3} usually entirely hyaline; a small apical spot. Anal streak present. Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale in basal third to half, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed, possibly with a minute apical notch.

(Description after White, 2006)