

***Dacus (Didacus) elatus* White, 2006**

Dacus (Didacus) elatus White, 2006: 89

Wing length, 6.7-7.2 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a long dark stripe (very tall spot). Frons, frontal setae 0-2, orbital setae 0-1 (all very reduced).

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for deep basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia (confined to anterior and distal margin areas); cell bm without microtrichia.

Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R_{4+5} (pale below R_{2+3}) before wing apex; apically expanded into a spot which reaches vein M. Anal streak present (colour extending beyond cell bcu). Cells bc and cell c slightly coloured (not as deep as costal band). Crossbanding; trace of mark at upper end of R-M; trace of a diffuse marking on DM-Cu.

Legs. Forefemur red-brown; mid- and hindfemora bicoloured (pale basal third, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)