

***Dacus (Didacus) fissuratus* White, 2006**

*Dacus (Didacus) fissuratus* White, 2006: 90

Wing length, 4.8 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a very tall dark spot on a dark face. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein  $R_{2+3}$  before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding; crossband on R-M, which continues as a diffuse mark across cell dm.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basal half, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. Aculeus with a minute apical notch; no torsion.

(Description after White, 2006)