

***Dacus (Didacus) frontalis* Becker**

*Dacus frontalis* Becker, 1922: 74

Wing length, 4.2-5.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep).

Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present or absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not or only slightly extending below vein  $R_{2+3}$  before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and hindfemora pale; midfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion; length, 1.3-1.7mm.

(Description after White, 2006)