

***Dacus (Didacus) jubatus* (Munro)**

Didacus jubatus Munro, 1984: 77

Wing length, 5.7 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta usually absent, at most a thin white seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; apically expanded into a spot which reaches about mid-depth of cell r_{4+5} . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline.

Without any crossbanding.

Legs. Fore- and midfemora indistinctly bicoloured (pale basally, red-brown apically); hindfemur pale.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Unknown.

Female. No available data [tip broken].

(Description after White, 2006)