

***Dacus (Didacus) nairobiensis* White, 2006**

Dacus (Didacus) nairobiensis White, 2006: 94

Wing length, 4.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 0, orbital setae 0.

Thorax. Scutum predominantly fulvous; postpronotal lobe yellow; notopleural callus concolorous with scutum; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum with a basal red-brown triangle. Anepisternum with a stripe from notopleural callus almost to katepisternum; not extended onto katepisternum. Laterotergal xanthine very small and confined to katatergite.

Thoracic setae. Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with microtrichia confined to anteriorly half. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; apically expanded into a spot which reaches nearly whole depth of cell r_{4+5} . Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding (except for a very slight infuscation of DM-Cu).

Legs. Femora pale coloured.

Abdomen. Predominantly fulvous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)