

***Dacus (Didacus) rugatus* Munro**

Lactodacus rugatus Munro, 1984: 96

Wing length, 5.4 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark stripe. Frons, frontal setae 2;

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep); sometimes almost concolorous with scutum. Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly deep, extending below vein R_{2+3} or to vein R_{4+5} , before wing apex; expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus blunt and with preapical "shoulder"; complete torsion.

(Description after White, 2006)