

***Dacus (Didacus) umbeluzinus* (Munro)**

Dixoodacus umbeluzinus Munro, 1984: 93

Wing length, 4.5-4.8 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; expanded into a small spot at apex. Anal streak variable (sometimes confined to within bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basally and red-brown apically); forefemur sometimes only darkened preapically.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus with a minute apical notch.

(Description after White, 2006)